



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NEBRASKA

Bravo WeatherStik®
For Control of Diseases on Long-Season Potatoes

EPA Reg. No. 50534-188-100
EPA SLN No. NE-090001

This label expires and must not be distributed or used in accordance with this SLN registration after September 1, 2020

KEEP OUT OF REACH OF CHILDREN
CAUTION

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

Crop	Disease (Pathogen)	Pints Product/A (lb ai/A)	Application Directions
Potato (Long-season varieties)	Late Blight (<i>Phytophthora infestans</i>)	Low rate: ¾ (0.6)	Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals (the minimum re-treatment interval is 5 days).
	Early Blight (<i>Alternaria solani</i>)	-then-	
	Botrytis vine rot (<i>Botrytis cinerea</i>)	Higher rates: 1 to 1½ (0.75 to 1.125)	Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur: <ul style="list-style-type: none"> • Vines close between rows • Late blight forecasting measures 18 disease severity values (DSV) • The crop reaches 300 P-days <p>Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe.</p> <p>Bravo WeatherStik may be applied</p>
	Blackdot (<i>Colletotrichum coccodes</i>)		

Crop	Disease (Pathogen)	Pints Product/A (lb ai/A)	Application Directions
			through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move or center pivot systems only). Do not exceed a 10 day interval between applications when using chemigation. See <i>Application and Calibration Techniques for Chemigation</i> on Bravo WeatherStik label
Specific Use Restrictions: Do not apply more than 21 ½ pints Bravo WeatherStik (16 lb ai) per acre on long-season varieties of potato during each growing season. Do not apply within 7 days of harvest.			

©2015 Syngenta

Bravo WeatherStik® and the Syngenta logo trademarks of a Syngenta Group Company

24(c) Sub-Registrant:
 Syngenta Crop Protection, LLC
 P. O. Box 18300
 Greensboro, NC 27419-8300

Label Code: NE0188019BA0515