



## Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NEBRASKA

**Bravo® Zn**  
**For Control of Diseases on Long-Season Potatoes**

**EPA Reg. No. 50534-204-100**  
**EPA SLN No. NE-090002**

**This label expires and must not be distributed or used in accordance with this SLN registration after September 1, 2020**

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING/AVISO**

### DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

Crop	Diseases (Pathogen)	Pints Product/A (lb ai/A)	Application Directions
<b>Potato</b> (Long-season varieties)	Late Blight ( <i>Phytophthora infestans</i> )	Low rate: 1 (0.6)	<p>Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals (the minimum re-treatment interval is 5 days).</p> <p>Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur:</p> <ul style="list-style-type: none"> <li>• Vines close between rows</li> <li>• Late blight forecasting measures 18 disease severity values (DSV)</li> <li>• The crop reaches 300 P-days</li> </ul> <p>Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe.</p> <p>Bravo Zn may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral</p>
	Early Blight ( <i>Alternaria solani</i> )	-then-	
	Botrytis vine rot ( <i>Botrytis cinerea</i> )	Higher rates: 1½ to 2 1/8 (0.75 to 1.125)	
	Black dot ( <i>Colletotrichum coccodes</i> )		

Crop	Diseases (Pathogen)	Pints Product/A (lb ai/A)	Application Directions
			move or center pivot systems only). Do not exceed a 10 day interval between applications when using chemigation. See <i>Application and Calibration Techniques for Chemigation</i> on Bravo Zn label.
<b>Specific Use Restrictions:</b> Do not apply more than 30½ pints Bravo Zn (16 lb ai) per acre on long-season varieties of potato during each growing season. Do not apply within 7 days of harvest.			

©2015 Syngenta

24(c) Sub-Registrant:  
 Syngenta Crop Protection, LLC  
 P. O. Box 18300  
 Greensboro, NC 27419-8300

Label Code: NE0204020BA0515