

FIFRA 24(c) Special Local Need Label (SLN)

For Distribution and Use Only in Nebraska

Control of Armyworms Infesting Dry Beans and Sugarbeet by Low Volume Aerial Application

Intrepid[®] 2F

EPA Reg. No. 62719-442

SLN NE-120001

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on the EPA registered label for EPA Reg. No. 62719-442.

Note: The use directions on this Special Local Need supplemental labeling supercede the aerial application use directions to apply in a minimum of 10 gallons per acre for the crops specifically listed on this supplemental labeling.

Refer to product label for Product Information, Mixing and Application directions.

Dry Beans

Application Rate (fl oz/acre)	Application Timing	Restrictions
8 - 16 (0.12 – 0.25 lb ai/acre)	Apply in a minimum of 5 gallons per acre. For mid- to late-season applications, heavier infestations, and under conditions in which thorough coverage is more difficult. For heavy infestations, continuous moth flights, and/or egg masses and larvae in all stages of development, a 7- to 14-day re-treatment interval is required to protect new growth until moth flights and/or larval infestations subside.	<ul style="list-style-type: none">• Preharvest Interval: Do not apply within 7 days of harvest.• Do not apply more than 16 fl oz per acre per application or more than a total of 64 fl oz of Intrepid 2F (1 lb ai) per acre per year.• Do not make more than 4 applications per acre per year.• Minimum Re-treatment Interval: 7 days• Do not use adjuvants in the tank mix when applying this product to dry beans.• See Rotational Crop Restrictions on the label affixed to the container.

Sugarbeet

Application Rate (fl oz/acre)	Application Timing	Restrictions
8 - 16 (0.12 – 0.25 lb ai/acre)	Apply in a minimum of 5 gallons per acre. Apply at egg hatch or when first signs of feeding occur. Use a higher rate for heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapply to protect new growth until moth flights and/or hits subside.	<ul style="list-style-type: none">• Preharvest Interval: Do not apply within 7 days of harvest.• Do not apply more than a total of 64 fl oz of Intrepid 2F (1 lb ai) per acre per year.• Minimum Re-treatment Interval: 14 days• See Rotational Crop Restrictions on the label affixed to the container.

Spray Adjuvants: The addition of agricultural adjuvants to sprays of Intrepid 2F may improve initial spray deposits, redistribution and weatherability. Select adjuvants that are recommended and registered for your specific use pattern and follow their use directions. When an adjuvant is to be used with this product, Dow AgroSciences

recommends the use of a Chemical Producers and Distributors Association certified adjuvant. Always add adjuvants last in the mixing process.

Spray volumes less than 10 gallons per acre may not achieve the same level of crop coverage as a 10 gallon volume. With reduced crop coverage, efficacy may be reduced.

Note: This product is toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

24(c) Registrant:

Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, Indiana 46268
1- 800-992-5994 (option #4)

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R846-086

Approved: __/__/__

Replaces: R846-086